


Fostera™ PCV MH

Description

For the vaccination of healthy pigs 3 weeks of age or older as an aid in preventing viremia, lymphoid depletion, and colonization of lymphoid tissue caused by porcine circovirus type 2 (PCV2), and as an aid in reducing PCV2 virus shedding and enzootic pneumonia caused by *Mycoplasma hyopneumoniae*. Duration of immunity studies are in progress.



Advantages

- The first and only combination vaccine to offer one-bottle, one-dose convenience with protocol flexibility!
- Built from the ground up – not a simple combination of two existing vaccines
- Uniquely developed to target the animal's immune response for protection against PCVAD caused by PCV2 and enzootic pneumonia caused by *M. hyopneumoniae*
- Aspects of human vaccine technology were applied to the manufacturing process
- The rigor put into every step of development, manufacturing, quality control, and testing means veterinarians and producers can trust Fostera PCV MH to help them achieve a high standard of consistent performance
- Developed by Zoetis with commitment to best practices and continuous quality improvements

Packaging

Fostera PCV MH is available in the following presentations:

- 50 dose – SKU #2381
- 250 dose – SKU #2382

Fosterera™ PCV MH

Dosage and Administration

Shake well.

Single dose: Administer a single 2-mL dose intramuscularly to pigs 3 weeks of age or older

Flexible dose: May also be given as two 1-mL doses administered 2 weeks apart to pigs 3 weeks of age or older

Precautions

Store at 2°–7°C (35°–45°F).

Do not freeze.

Use entire contents when first opened.

Do not vaccinate within 21 days before slaughter.

Contains gentamicin and thimerosal as preservatives.

Allergic reactions, although rare, may occur.

The antidote is epinephrine.

Contact Information

Technical inquiries should be directed to Zoetis Inc. Veterinary Services:
(888) 963-8471 (USA).

For additional information, contact your local Zoetis representative.